

Dynamical turbulent flow on the Galton board with friction

A. D. Chepelianskii^(a) and D. L. Shepelyansky^(b)

^(a) *Lycée Pierre de Fermat, Parvis des Jacobins, 31068 Toulouse Cedex 7, France*

^(b) *Laboratoire de Physique Quantique, UMR 5626 du CNRS, Université Paul Sabatier, 31062 Toulouse Cedex 4, France*
(January 1, 2001)

We study numerically and analytically the dynamics of particles on the Galton board, a regular lattice of disc scatters, in the presence of a constant external force and friction. It is shown that under certain conditions friction leads to the appearance of a strange chaotic attractor in an initially conservative Hamiltonian system. In this regime the particle flow becomes turbulent and its average velocity depends in nontrivial manner on friction and other system parameters. We discuss the applications of these results to the transport properties of suspended particles in a laminar viscous flow streaming through scatters.

PACS numbers: 05.45.Ac, 47.52.+j, 72.20.Ht

It is well known that dissipation can lead to the appearance of strange chaotic attractors in nonlinear nonautonomous dynamical systems [1,2]. In this case the energy dissipation is compensated by an external energy flow so that stationary chaotic oscillations set in on the attractor. Such an energy flow is absent in the Hamiltonian conservative systems and therefore the introduction of dissipation or friction is expected to drive the system to simple fixed points in the phase space. This rather general expectation is surly true if the system phase space is bounded. However a much richer situation appears in the case of unbounded space, where unexpectedly a strange attractor can be induced by dissipation in an originally conservative system. To investigate this situation we study the dynamics of particles on the Galton board in the presence of a constant external field and friction. This board, introduced by Galton in 1889 [3], represents a triangular lattice of rigid discs with which particles collide elastically. For the case of free particle motion, the collisions with discs make the dynamics completely chaotic on the energy surface as it was shown by Sinai [4]. In this paper we study how the dynamics is changed if the particles are affected by a friction force $\mathbf{F}_f$ which is assumed to be proportional to the particle velocity $\mathbf{v}$ and directed against it: $\mathbf{F}_f = -\gamma\mathbf{v}$. In the free space without discs, an external field with an effective force $\mathbf{f}$ creates a stationary particle flow with the velocity $\mathbf{v}_f = \mathbf{f}/\gamma$. All perturbations decay to this flow with the rate proportional to γ so that this laminar flow can be considered as a simple attractor. It is interesting to understand how this flow will be affected by the scattering on regularly placed discs. As we will show latter on, in this case the flow can become turbulent and chaotic, and demonstrate many unusual properties (see Fig.1). The Hamiltonian dynamics of the model has been studied in great detail (see [5] and Refs. therein) however, as for our knowledge, the effects of friction were not addressed up to now. The investigation of the dynamics in this regime is interesting from the viewpoint of general phys-

ical grounds. But we also note that physically such a situation can correspond to the motion of small particles in a viscous liquid flow which laminarly streams through large scatters that in our case have the form of discs. The same model can describe the electron dynamics in antidot superlattice which has been experimentally realized in semiconductor heterostructures [6]. In such structures the effects of classical chaos play an important role [7] and the effects of friction we discuss here can appear for relatively strong electric fields.




FIG. 1. Chaotic flow on the Galton board. Here the distance between discs is $R = 2.24$, $\mathbf{f} = (-0.5, -0.5)$ and $\gamma = 0.1$. Initially 100 particles are distributed homogeneously along a straight line segment in the upper right corner, their color homogeneously changes from red to green along this segment.

To study the dynamics of this model, we fix the disc radius $a = 1$, and the mass of the particles $m = 1$ so that the system is characterized only by three parameters: the distance R between discs packed into equilateral

triangles all over the (x, y) plane; the friction coefficient γ and the external force of strength f . For $R \leq R_c = 4/\sqrt{3}$ there is no straight line, by which a particle can cross the whole plane without collisions (at $f = 0, \gamma = 0$), and we start the discussion from this case. The particle dynamics is simulated numerically by using the exact solution of Newton equations between collisions, and by determining the collision points with the Newton algorithm. This way the trajectories are computed with high precision, and for example for $\gamma = 0$ the total energy is conserved with a relative precision of 10^{-14} . A typical example of the chaotic flow formed by an ensemble of particle trajectories is shown in Fig. 1. To illustrate the mixing properties of the flow we attributed a color to each trajectory that allows to follow their interpenetration and spreading. This figure shows that there is a certain penetration depth of one color into another, however this depth is finite since on average the initial color repartition is still visible.




FIG. 2. Example of a single trajectory for the case of Fig. 1 shown on a small (main figure) and large (inset) scales. The drift velocity of the flow is $v_f \approx 0.13$.

The properties of color penetration can be understood from the analysis of single trajectory dynamics. Such a typical example is presented in Fig. 2. It shows that trajectory flows with an average constant velocity v_f in the direction of the external force. This drift velocity is constant only on average since on a smaller scale the particle moves chaotically between scatters (see Fig. 2). Even if the average flow is regular, on a small scale the particle dynamics is chaotic and since the dynamics is dissipative this means that the motion develops on a strange chaotic attractor. This attractor is induced by dissipation in a conservative Hamiltonian system. For the case of Fig. 2 the drift velocity is relatively large and the particle does not have enough time to move around many scatters in the direction perpendicular to the flow. As

a result the penetration depth for color mixing is not very large. However for smaller friction, the drift velocity becomes smaller and the penetration depth increases so that the particle makes many turns around discs as it is shown in Fig. 3. Surprisingly, the drift velocity v_f *increases* with the growth of the friction coefficient γ . This dependence is opposite to the case without scatters where v_f drops with the γ growth.




FIG. 3. Single trajectory for the same parameters as in Figs. 1,2 but with smaller friction $\gamma = 0.004$; on a large scale the particle moves with a drift velocity $v_f \approx 0.05$ in the direction of external force.




FIG. 4. Dependence of the flow velocity v_f on the friction coefficient γ and external field strength f for $R = 2.24$, $\mathbf{f}/|\mathbf{f}| = (-1, -1)$ and $0.001 \leq \gamma \leq 0.4$; $0.5 \leq f \leq 32$. Numerical results are shown by circles, the straight line shows the dependence $v_f \propto (\gamma f)^{1/3}$. Inset shows the dependence of v_f on $\Delta R = R - 2$ for $\gamma = 0.1$ and $\mathbf{f} = (-0.5, -0.5)$: points give numerical data, lines show the slopes 0.5 and 1.

The dependence of the drift velocity v_f on γ and on the external force $f = |\mathbf{f}|$ is obtained by averaging over many trajectories. Each trajectory is followed for a sufficiently long time so that the distance from the origin becomes

larger than a few thousands units. Up to fluctuations, the flow is always directed along the applied force $\mathbf{f}$. The resulting dependence is shown in Fig. 4 according to which $v_f \propto (\gamma f)^{1/3}$. To understand the origin of this unexpected dependence on parameters we also compute the average velocity square $v^2 = \langle v_x^2 + v_y^2 \rangle$ and determine how it varies with f and γ . The data presented in Fig. 5 give a dependence of the form: $v^2 \propto f^{4/3}/\gamma^{2/3}$. We stress that v_f and v^2 are independent of initial velocity of the particle.

To explain the observed scaling relations for the dynamical turbulent flow on the Galton board we give the following theoretical arguments. In the regime with weak friction the particles start to diffuse among discs in a chaotic manner with the diffusion rate $D = vl/2$ where v is the velocity of the particle and l is the mean free path [8]. For $R \sim 1$ we have $l \sim R \sim 1$ while the dependence of l on R will be discussed in more detail latter. During the dissipative time scale $\tau_\gamma = m/\gamma$ this diffusion leads to the particle displacement $\Delta r \sim \sqrt{Dm/\gamma}$ along the direction of applied field. Here, for generality we introduced the particle mass m . This gives the change in the potential energy $U \sim f\Delta r \sim f\sqrt{Dm/\gamma}$. In the stationary regime at time $t \gg \tau_\gamma$, this potential energy should be comparable with the kinetic energy of the particle so that $U \sim mv^2$. As a result we find that the average velocity square is

$$v^2 \sim f^{4/3}(l/\gamma m)^{2/3}, \quad (1)$$

that is in agreement with the data of Fig. 5.




FIG. 5. Dependence of average v^2 on f and γ for $R = 2.24$, $\mathbf{f}/|\mathbf{f}| = (-1, -1)$ and parameter range $0.001 \leq \gamma \leq 0.4$; $0.5 \leq f \leq 32$. Circles show numerical data, the straight line gives the slope 1.

The relation (1) allows to determine the mobility μ which relates the velocity of the flow with the applied force: $\mathbf{v}_f = \mu \mathbf{f}$. Indeed during the time τ_c between collisions the particle is accelerated by force that gives the average drift velocity $v_f = f\tau_c/m$ and $\mu = \tau_c/m$.

Since the dynamics is chaotic the direction of velocity is changed randomly after each collision so that v_f is accumulated only between collisions. The time τ_c is determined by the mean free path l and the average velocity v : $\tau_c = l/v = 2D/v^2$ and $\mu = D/(mv^2/2)$. In fact this is the Einstein relation according to which the mobility is given by ratio of the diffusion rate to the average kinetic energy (temperature) [9]. Together with (1) this gives the drift velocity of the chaotic flow:

$$v_f \sim l^{2/3}(f\gamma/m^2)^{1/3}, \quad (2)$$

which is in a good agreement with the numerical results in Fig. 4. On average the particle moves with the velocity v_f in the direction of the force (see an example in the inset of Fig. 2). The amplitude of fluctuations around this direction is $\Delta r \sim \sqrt{Dm/\gamma}$ which also determines the color mixing depth (see Fig. 1).

The relations (1), (2) allow to estimate the value of the Lyapunov exponent λ . Indeed, the particle moves with a typical velocity v and as in the case of the Sinai billiard with $l \sim R \sim 1$ we have $\lambda \sim v/l$. Therefore in the regime when:

$$\gamma < \gamma_c \sim \sqrt{f m / l}, \quad (3)$$

the value of λ is much larger than the dissipation rate γ/m . As a result for $\gamma \ll \gamma_c$ the strange attractor is fat and its fractal dimension is close to the maximal dimension 4, which is determined by the number of degrees of freedom (we remind that contrary to the nondissipative case the energy is not conserved).

The physical meaning of the critical γ_c border is the following: for $\gamma \gg \gamma_c$ the dissipation time τ_γ becomes much shorter than the time between collisions τ_c . In this case the dissipation dominates chaos and the strange attractor degenerates into a simple attractor. Among such simple attractors we observed trajectories that move regularly under some angle to the applied force but more often trajectories that simply drop to a fixed point on a disc where the applied force is perpendicular to the disc border. Of course the numerical simulations can not guarantee that the chaotic attractor at $\gamma < \gamma_c$ will not degenerate to a simple one at very long times. However all our simulations performed with high computer precision give no indications in this direction even when we followed trajectories up to very far distances from the origin (bigger than 10^5 length units).

The equation (3), which determines the border for disappearance of chaotic flow, is written for the case $l \sim R \sim 1$. Therefore, it should be modified when $R \gg 1$ and $\Delta R = R - 2 \ll 1$. Indeed for $R \gg 1$ the mean free path l is given by $l = 1/(\sigma n)$ where $\sigma = 2a = 2$ is the disc cross section and $n = 2/(\sqrt{3}R^2)$ is the density of scatters. Then the time between collisions is l/v and from equation (1) we obtain $\gamma_c \sim mv/l \sim \sqrt{fma}/R$. In the other limit $\Delta R \ll 1$ we may assume that $D \sim v\Delta R$,

$l \sim \Delta R$ and the escape time from a cell between three discs is $\tau_{esc} \sim a^2/D$. Therefore the dissipation becomes dominant and the chaotic attractor disappears at $\gamma > \gamma_c \sim m/\tau_{esc} \sim \sqrt{mf/a^3} \Delta R$.

The above changes in the mean free path l also affect the drift velocity of the flow through the relation (2). Indeed for $\Delta R \ll 1$ we expect $l \sim \Delta R$ that gives $v_f \propto \Delta R^{2/3}$. This dependence is close to the numerical data shown in Fig. 4 (inset) even if the numerical value of the exponent is approximated better by 0.5. In the other limit $R \gg 1$, we have $l \sim R^2/a$ that gives $v_f \propto R^{4/3}$. This power dependence is in a satisfactory agreement with our numerical data (inset in Fig. 4) although the numerical value of the exponent is closer to 1. We attribute these small deviations in the exponent values to a restricted interval of variation in R . Actually we can not use very large/small values of ΔR since in these limits the value of γ becomes comparable with γ_c and chaotic attractor disappears. We note that according to our data (see Fig. 4) the strange attractor exists even in the case $R > R_c = 4/\sqrt{3}$ when at $f = 0, \gamma = 0$ there are straight trajectories crossing the whole plane without collision. Apparently the contribution of these orbits is not significant if $\gamma > 0$ and $\mathbf{f}$ is not directed along these lines.

Let us now discuss a few consequences of the obtained results. The introduction of scatters qualitatively changes the flow properties: instead of a laminar flow we obtain a turbulent/chaotic one. The flow rate v_f depends on the applied force f and γ in a nonlinear way given by equation (2). Contrary to the laminar case where $v_f = f/\gamma$ in presence of scatters the flow velocity depends also on the mass of particles m . In fact light particles stream with higher velocity. This property of chaotic flow can be used for mass separation of particles suspended in viscous flows, for example isotopes. It is also interesting to note that in the derivation of equations (1), (2), (3) we didn't use any specific properties of the disc distribution in the plane. Therefore these results remain also valid for a random distribution of disc scatters in the plane. We wonder if some rigorous mathematical results can be obtained in the limit of weak friction to prove the existence of a chaotic attractor for a regular or random distribution of discs.

We think that the obtained results represent certain interest for transport properties of particles suspended in a viscous flow streaming through a system of large scatters. Indeed, when a liquid streams laminarly through scatters with the velocity v_s the flow of suspended particles is characterized by the relations (1), (2), (3) with $f_{eff} = v_s \gamma_{eff}$ where γ_{eff} is an effective friction created by the viscosity of the liquid and f_{eff} is the resulting effective force acting on the suspended particles. Then for weak friction the drift velocity of particles v_f is smaller than the velocity of the liquid v_s . Such kind of transport

can be studied experimentally with viscous liquids and its investigation can contribute to a better understanding of the interplay between dissipation, turbulence and chaos.

-
- [1] A. Lichtenberg and M. Lieberman, *Regular and chaotic dynamics*, Springer, Berlin (1992).
 - [2] E. Ott, *Chaos in dynamical systems*, Cambridge Univ. Press (1993).
 - [3] F. Galton, *Natural inheritance*, London (1889).
 - [4] I. P. Kornfeld, S. V. Fomin, and Ya. G. Sinai, *Ergodic theory*, Springer, Berlin (1982).
 - [5] P. Gaspard and G. Nicolis, Phys. Rev. Lett. **65**, 1693 (1990); H. van Beijeren and J. R. Dorfman, Phys. Rev. Lett. **74**, 4412 (1995).
 - [6] D. Weiss, M. L. Roukes, A. Mesching, P. Grambow, K. von Klitzing, and G. Weiman, Phys. Rev. Lett. **66**, 2790 (1991).
 - [7] R. Fleischmann, T. Geisel, and R. Ketzmerick, Phys. Rev. Lett. **68**, 1367 (1992).
 - [8] J. M. Ziman, *Principles of the theory of solids*, Cambridge Univ. Press (1972).
 - [9] L. D. Landau and E. M. Lifshitz, *Hydrodynamics*, Nauka, Moscow (1986).